

Early Real-time Estimation of Infectious Disease Reproduction Number

Bahman Davoudi¹, Babak Pourbohloul^{1,2,*}, Joel C. Miller³, Rafael Meza¹,
Lauren Ancel Meyers⁴, David J. D. Earn⁵

October 1, 2009

Abstract

When an infectious disease strikes a population, the number of newly reported cases is often the only available information that one can obtain during early stages of the outbreak. An important goal of early outbreak analysis is to obtain a reliable estimate for the basic reproduction number, R_0 , from the limited information available. We present a novel method that enables us to make a reliable real-time estimate of the reproduction number at a much earlier stage compared to other available methods. Our method takes into account the possibility that a disease has a wide distribution of infectious period and that the degree distribution of the contact network is heterogeneous. We validate our analytical framework with numerical simulations.

¹ Division of Mathematical Modeling, British Columbia Centre for Disease Control,
Vancouver, British Columbia, Canada

² School of Population & Public Health, University of British Columbia, Vancouver, British
Columbia, Canada

³ Department of Epidemiology, Harvard School of Public Health, Boston, Massachusetts,
U.S.A.

⁴ Section of Integrative Biology, Institute for Cellular and Molecular Biology, University of
Texas at Austin, Austin, Texas, U.S.A.

⁵ Department of Mathematics & Statistics, McMaster University, Hamilton, Ontario, Canada

**Corresponding Author*

1 Introduction

The basic reproduction number, R_0 , is a fundamental characteristic of an infectious disease. It is generally defined as the expected number of new infections caused by a typical individual during the whole period of his/her infection, in a totally susceptible population (Anderson & May, 1991; Anderson & Britton, 2000; Dietz, 1993). Because R_0 is a simple scalar quantity, and perhaps because in many circumstances it determines the expected final size of an outbreak (Kermack & McKendrick, 1927; Dietz, 1993; Ma & Earn, 2006; Arino *et al.*, 2007; Brauer, 2008) it has been widely used as a gold standard to gauge the degree of threat that a specific infectious agent will pose as an outbreak progresses (Hethcote, 2000; Lipsitch *et al.*, 2003; Fraser *et al.*, 2009). While it is clear that knowing the value of R_0 can be very useful to policy-makers who are responding to an infectious disease invasion, it is not straightforward to obtain a reliable estimate of R_0 , especially in early stages of an outbreak before large scale uncontrolled transmission has taken place and before the basic biology and transmission pathways of the pathogen have been characterized.

Early in an outbreak, the pattern of disease spread is predominantly under the influence of the probabilistic nature of infection transmission. Consequently, a wide array of ultimate outcomes is possible, ranging from the outbreak fizzling out even in the absence of intervention to circumstances where the initial stage expands into a large-scale epidemic. Once a full-blown epidemic develops, several assumptions can be made that simplify the estimation of R_0 , as has been discussed in detail in the literature (Anderson & May, 1991; Dietz, 1993; Hethcote, 2000; Ma & Earn, 2006). In many circumstances, however, it is crucial to assess the impact of various intervention strategies before a large-scale epidemic occurs. In doing so, stochastic manifestations of disease transmission as well as the underlying structure of the contact network

should be taken into account. The first aspect has been widely studied during the past. For example, the Reed-Frost model is a particular case of a chain-binomial stochastic model, where each infected individual can infect susceptible individuals independently, and those individuals are assumed to have the same contact rate with one another (Abbey, 1952; Addy *et al.*, 1991; Ball, 1986). Another example is the methodology developed by Becker (Becker, 1976) and Ball and Donnelly (Ball & Donnelly, 1995) based on a branching process susceptible-infected-recovered (SIR) model. Branching processes have received wide attention because they facilitate the evaluation of the reproduction number as well as the final epidemic size and epidemic probability (Guttorp, 1991). More recently, the 2003 global outbreak of severe acute respiratory syndrome (SARS) inspired the development of new methodologies based on knowledge of the daily number of new cases and the distribution of the serial interval between successive infections (Wallinga & Teunis, 2004; Cauchemez *et al.*, 2006*a,b*). Unfortunately, none of these methods take into consideration the underlying contact network behind the epidemic process.

In this paper, we propose a new methodology to estimate the basic reproduction number together with other useful epidemiological quantities at an early stage of an outbreak, while accounting for both the stochastic nature of the transmission process and the underlying structure of the contact network. The main parameters used in this approach, i.e., the quantities we require in order to apply the method, are the *infection rate probability*, the *removal rate probability* and the *number of newly-infected* individuals per unit of time (case-notification data). The infection rate quantifies crucial information about the biology of the disease. For instance, for a viral infection, the infection rate, in principle, may be related to the viral load of infectious individuals. The removal rate indicates how quickly the infectious individuals are removed from disease dynamics. This rate can be related to death or various means of interven-

tion such as quarantine, reduction of social activity due to severity of illness, behavior change, or other factors. The product of infection rate probability and the cumulative removal rate probability is related to the *generation interval distribution* (Wallinga & Lipsitch, 2007) (see section VI). The three main parameters will be discussed in more detail throughout the paper. We will derive a stochastic renewal equation that allows us to understand how the number of newly-infected individuals is related to the preceding number of infected cases. Unlike existing approaches (e.g., Wallinga and Teunis (Wallinga & Teunis, 2004)), our method incorporates the degree distribution of the social contact network. We will show how knowledge of the social contact network can be utilized to obtain a more reliable estimate of the reproduction number.

This paper is organized as follows. In Section 2 we introduce the ideas of contact network epidemiology and define the concepts of infectious rate, recovery rate and transmissibility. In Section 3 we show some examples of the spread of an infectious agent on a contact network. In Section 4 we outline a general framework to estimate the reproductive number assuming that all information about a specific realization of the epidemic process up to time t is known. In Section 5 we derive several expressions that allow us to estimate the basic reproduction number, R_0 , at an early stage of an outbreak and show some examples of the methodology. Finally, in section 6 we show the relationship between our results and the methodology developed by Wallinga and Lipsitch (Wallinga & Lipsitch, 2007), which is a special case of the framework presented in this paper.

2 Network basis

In this section we introduce briefly the ideas of contact network epidemiology and define the concepts of infectious rate, recovery rate and transmissibility. We map a collection of N individ-

uals to a network in which each vertex represents an individual and each edge shows a pathway of possible infection transmission between two individuals. The degree of each individual, i.e., the number of contacts that s/he has - or the number of edges emanating from a vertex - is denoted by κ , and p_κ denotes the degree distribution, i.e., the probability that a randomly chosen vertex has degree k (k contacts). Several important quantities can be derived once a network's degree distribution is known. All moments of the degree distribution can be obtained since $\langle k^n \rangle = \sum_{\kappa=0}^{\infty} k^\kappa p_\kappa$. For $n = 1$, $\langle k \rangle$ is the average number of nearest neighbors of a randomly chosen individual, which we denote z_1 . The average number of second degree neighbors of a randomly selected individual, z_2 , can be expressed as $\langle k^2 \rangle - \langle k \rangle$ (Newman, 2002). To estimate R_0 , we count the number of edges along which an individual can infect others, once that individual has become infected. This quantity, the *excess degree*, is the number of edges emanating from a vertex (individual), excluding the edge that was the source of the focal individual's infection. In fact, the *average excess degree* is simply the ratio $\frac{z_2}{z_1}$ (Noel *et al.*, 2009).

We denote the time at which an individual acquires the infection, by t_I , the time since acquiring infection by $\tau = t - t_I$ (also known as age of infection) and the total time that the individual can cause infection (time-to-removal) by t_R . While harboring the infection, the person is first *exposed* (i.e., infected but not infectious) and then *infectious* (either symptomatically or asymptotically). The individual may also *recover*, by which we mean only that s/he can no longer transmit the infection, not that s/he has necessarily completely cleared the pathogen; for some diseases, after a temporary recovery, the person might become infectious again. Knowing that an individual acquires infection at a given time t_I , various states of infectiousness of this individual can be encapsulated within the hazard of infection or *infectivity function*, $\lambda_i(\tau)$. The infectivity function measures the instantaneous risk of disease transmission across an edge.

This means that the conditional probability that infection occurs across an edge between times τ and $\tau + d\tau$ given it did not occurred by time τ is equal to $\lambda_i(\tau)d\tau$. Typically $\lambda_i(\tau)$ is zero initially (i.e., during the latent period), increases to a certain level and then declines (during the infectious period) before finally vanishing and remaining zero (i.e., in the case of permanent recovery or death). Figure 2 shows four hypothetical infectivity functions, the first of which is the typical case. In practice, the functional form of $\lambda_i(\tau)$ should be estimated from the actual transmission profile corresponding to a specific disease. Note that because of the limited restrictions on the functional form of $\lambda_i(\tau)$, must be a non-negative integrable function, the methodology we present here applies to any staged progression transmission process, such as an $SE_1E_2\dots E_mI_1I_2\dots I_nR$ process, where S , E_i and I_j ($i = 1, \dots, m$ and $j = 1, \dots, n$) and R are the susceptible, exposed (infected but not infectious), infectious and recovered classes, respectively.

Using the infectivity function, we can evaluate the probability that transmission occurs across an edge during a specific time period. In particular, given the removal time t_R of an individual, the transmissibility $T(\tau, t_R)$ - the probability that the individual transmitted the disease to one of its neighbors before τ units of time (since acquiring the infection) satisfies (Cox & Oakes, 1984; Newman, 2002):

$$\left\{ \begin{array}{ll} T(\tau, t_R) = 1 - \exp \left(- \int_0^\tau \lambda_i(u) du \right) & \tau < t_R \\ T(\tau, t_R) = T(\infty, t_R) = 1 - \exp \left(- \int_0^{t_R} \lambda_i(u) du \right) & otherwise \end{array} \right\}. \quad (1)$$

Let $\phi(\tau, t_R)$ denote the logarithmic derivative of $T(\tau, t_R)$, which we will refer to as the *infectivity rate* i.e., let

$$\phi(\tau, t_R) = \frac{1}{T(\tau, t_R)} \frac{dT(\tau, t_R)}{d\tau}. \quad (2)$$

Then it follows that $T(\tau, t_R)$ can also be expressed as:

$$T(\tau, t_R) = \exp \left(\int_0^\tau \phi(u, t_R) du \right). \quad (3)$$

In general, the time-to-removal period, t_R , varies from one individual to another and there is no *a priori* knowledge of the exact value of this quantity for each individual. Therefore we must account for its variability as well. Let $\lambda_r(t_R)$ denote the *removal function* or recovery hazard, i.e., the instantaneous rate of recovery for an individual t_R units of time after acquiring the infection. This means that the conditional probability that an individual is removed between times t_R and $t_R + dt_R$ given it was not removed by time t_R is equal to $\lambda_r(t_R)dt_R$. Let $\Psi(t_R)$ denote the probability that an individual has time-to-removal period $\geq t_R$, then (Cox & Oakes, 1984)

$$\Psi(t_R) = \exp \left(- \int_0^{t_R} \lambda_r(u) du \right), \quad (4)$$

subject to the condition $\Psi(\infty) = 0$. We define the probability density function $\psi(t_R) = -\frac{d\Psi(t_R)}{dt_R}$ (or $\Psi(t_R) = \int_{t_R}^\infty \psi(t_R) dt_R$).

Using equation (1) and $\psi(t_R)$ one can calculate the expected value of the transmissibility across an edge τ units of time after infection as:

$$T(\tau) = \int_0^\infty \psi(t_R) T(\tau, t_R) dt_R. \quad (5)$$

Finally, we define the overall expected transmissibility, i.e., the transmissibility when we wait for an adequately long time ($t \rightarrow \infty$), as

$$T = T(\infty) = \int_0^\infty \psi(t_R) T(\infty, t_R) dt_R. \quad (6)$$




Figure 1: Four hypothetical profiles illustrating various infectivity functions, $\lambda_i(\tau)$.

The reproduction number can now be expressed as $R_0 = \frac{z_2}{z_1}T$, and represents the average number of infections caused by an infected individual before it is removed.

3 Disease dynamics on networks

In this section we present some examples of the spread of an infectious agent on a contact network. The pattern of disease spread on a network can be categorized by three different regimes: stochastic, exponential, and decline. The process of disease spread is indeed stochastic in nature, given that the disease transmission along an edge occurs in a probabilistic manner, and that the degree of each newly infected individual cannot be determined *a priori*. The stochastic behavior is dominant in the initial state of disease spread when the number of infectious individuals is comparatively small (stochastic regime). The effect of stochasticity becomes much less relevant when the number of newly infected individuals dramatically increases, as in this case where the stochastic fluctuations (noise) are overwhelmed by a smooth “signal” (exponential regime). In this regime, the number of new infections occurring at time t , $J(t)$, grows




Figure 2: Two hypothetical realizations of an epidemic process on a network. The left, middle and right hand sides represent stochastic, exponential and decline regimes, respectively.

exponentially and therefore can be expressed as:

$$J(t) \simeq \exp(\alpha t). \quad (7)$$

Progression of disease spread starts to decline as the total number of infected cases becomes comparable to the size of network. This is when the finite-size effects become important (Noel *et al.*, 2009) (declining regime). All three regimes are represented in figure 2.

Figures 3 and 4 show two examples of the epidemic behavior in the three regimes. The algorithm used to simulate the spread of an infectious agent on a contact network is described in the appendix. The left panel of figure 3 shows two epidemic events unfolding on a binomial network $p_k = \binom{N}{k} p^k (1-p)^{N-k}$ with $z_1 = 5$ (or $p = z_1/(N-1)$), $N = 100,000$ nodes and $\{\lambda_i = 0.12771, \lambda_r = 0.25\}$. We use the tilde notation to make the distinction between the realization of a stochastic process (with tilde) and its expected (average) value (without tilde). Using equation (3) the ultimate transmissibility can be calculated as $T = 0.32727$. The stochastic, exponential and decline regimes are approximately between $0 \leq t < 20$, $20 \leq t < 40$




Figure 3: The number of newly-infected cases (left panel) and its logarithm (right panel) for a binomial network with $z_1 = 5$, $\lambda_i = 0.12771$ and $\lambda_r = 0.25$. Two independent epidemic realizations are shown in each panel (green and blue).

and $40 \geq t$ time units, respectively. Also, in the right panel of figure 3, we show the variation of $\ln(\tilde{J}(t))$ over time. The two solid lines $y_1, y_2 \approx \alpha t$ with $\alpha_1 = \alpha_2 = 0.26$ are the tangents to the curves in the exponential regime. Figure 4 shows the same results for an exponential network with $p_k = (1 - e^{-1/\kappa}) e^{-k/\kappa}$ and $\kappa = 4$ ($z_2/z_1 = 7.0416$). In this example, the stochastic, exponential and declining regimes are approximately between $0 \leq t < 10$, $10 \leq t < 20$ and $20 \leq t$ time steps, respectively. The lines $y_1, y_2 \approx \alpha t$ with $\alpha_1 = \alpha_2 = 0.516$ are the tangents to the curves in the exponential regime. Here, we used epidemic simulation outputs to generate “synthetic” time series data. In practice, α can be estimated from real-life time series data if the outbreak progresses beyond the stochastic regime. As can be seen in figures 3 and 4, the number of newly infected cases at time t , $\tilde{J}(t)$, generally resembles a noisy Gaussian-type function if the pattern of disease spread has a chance to grow substantially.

4 Stochastic dynamics of disease

In this section we outline a general framework to estimate the reproductive number assuming that all information about a specific realization of the epidemic process up to time t is known.




Figure 4: The number of newly-infected cases (left panel) and its logarithm (right panel) for an exponential network with $\kappa = 4$, $\lambda_i = 0.12771$ and $\lambda_r = 0.25$. Two independent epidemic realizations are shown in each panel (green and blue).

Let $\tilde{\Psi}(t - \tau, t)\tilde{J}(t - \tau)$ denote the fraction of individuals infected at time $t - \tau$ that remain infectious at time t . As mentioned before, the tilde is used to denote that a quantity corresponds to a specific realization of the epidemic process. Let $\tilde{Z}(t - \tau, t)$ denote the average (per individual) excess degree at time t . The aggregated excess degree (total number of links) of the individuals *infected* at time $t - \tau$ that are still infectious at time t is then given by $\tilde{J}(t - \tau)\tilde{\Psi}(t - \tau, t)\tilde{Z}(t - \tau, t)$. Let $\tilde{T}(t, \tau, t_R \geq \tau)\phi(\tau, t_R \geq \tau)d\tau$ denote the fraction of these links that actually transmit the disease at time t (*infectious*). Then the number of new infections occurring at time t satisfies the renewal equation:

$$\tilde{J}(t) = \int_0^t \tilde{J}(t - \tau)\tilde{\Psi}(t - \tau, t)\tilde{Z}(t - \tau, t)\phi(\tau, t_R \geq \tau)\tilde{T}(t, \tau, t_R \geq \tau)d\tau. \quad (8)$$

$\tilde{J}(t)$ is the only dependent variable in the above formula.

As an outbreak progresses, at any given time there is a population of infected individuals, $\tilde{I}(t)$, and a population of removed individuals, $\tilde{R}(t)$. The number of *affected* individuals - the total number of infected individuals at time t and those who recovered/removed by time t - is given by




Figure 5: This figure illustrates the dependency of $J(t)$ on its past values. Only a fraction of the cases infected at former time $t - \tau$, $\tilde{J}(t - \tau)$, contributes to the infection at the current time t , $\tilde{J}^i(t, \tau) = \tilde{J}(t - \tau)\Psi(t - \tau, t)$, while the rest, $\tilde{J}^r(t, \tau) = \tilde{J}(t - \tau)[1 - \Psi(t - \tau, t)]$, have already been removed.

$$\tilde{\mathcal{I}}(t) + \tilde{\mathcal{R}}(t) = \int_0^t \tilde{J}(t - \tau) d\tau \quad (9)$$

where the total number of infected cases is given by

$$\tilde{\mathcal{I}}(t) = \int_0^t \tilde{J}(t - \tau) \tilde{\Psi}(t - \tau, t) d\tau \quad (10)$$

which in turn, implies that

$$\tilde{\mathcal{R}}(t) = \int_0^t \tilde{J}(t - \tau) [1 - \tilde{\Psi}(t - \tau, t)] d\tau \quad (11)$$

The total excess degree of *removed* individuals is given by

$$\tilde{Z}^r(t) = \int_0^t \tilde{J}(t - \tau) \tilde{Z}(t - \tau, t) [1 - \tilde{\Psi}(t - \tau, t)] d\tau \quad (12)$$

It is worth emphasizing that $\tilde{Z}^r(t)$ is the total number of edges that could have transmitted the infection by already removed individuals by time t , while only a fraction of them actually

transmitted the disease successfully. This latter fraction is given by $\tilde{\mathcal{I}}^r(t) + \tilde{\mathcal{R}}^r(t)$, where $\tilde{\mathcal{I}}^r$ and $\tilde{\mathcal{R}}^r$ are the number of infectious and removed individuals at time t , infected by individuals who are already removed at time t . In the same manner, one can define $\tilde{\mathcal{I}}^i(t)$ and $\tilde{\mathcal{R}}^i(t)$ as the number of infectious and removed individuals who acquired infection from those who remain themselves infectious at time t . Fig. 6 illustrates how individuals in various states infect each other, where the arrows show “who is infected from whom”. More precisely, the arrows establish the causality link by showing where the superscript index (the predecessor) may come from. For instance, a recovered/removed predecessor, denoted by superscript r in $\tilde{\mathcal{R}}^r(t)$, can only come from a group whose members were recovered while their predecessors were still infectious ($\tilde{\mathcal{R}}^i$), or from a group in which the person and his/her predecessor are removed (denoted by the self-loop from $\tilde{\mathcal{R}}^r$ to itself). Similarly, the recovered predecessor (r) of an infected person ($\tilde{\mathcal{I}}^r(t)$), may only come from a group of recovered individuals whose respective predecessors were infectious ($\tilde{\mathcal{R}}^i(t)$) or from a group whose respective predecessors were already recovered ($\tilde{\mathcal{R}}^r(t)$). The expected *transmissibility of the removed individuals* $\bar{T}^r(t)$ at time t can thus be defined as

$$\bar{T}^r(t) = \frac{\tilde{\mathcal{I}}^r(t) + \tilde{\mathcal{R}}^r(t)}{\tilde{\mathcal{Z}}^r(t)}. \quad (13)$$

It is not possible to obtain a closed form equation for $\tilde{\mathcal{I}}^r$ and $\tilde{\mathcal{R}}^r$ in terms of the above-mentioned quantities, because the information about who gets infected from whom is not included in equation (8). However, estimates for the expected values for these quantities can be calculated and are given in the next section. This will allow us to estimate the basic reproduction number R_0 .




Figure 6: This figure illustrates various options of causality between infected and recovered cases. Letter (I or R) denotes the current status of an individual, and the superscript (i or r) correspond to the status of the person who passed on the infection to this individual. $I^r(t) + R^r(t)$ constitutes the number infections caused by the already removed individuals.

5 Quasi-stochastic dynamics of disease spread

In this section we derive several expressions that allow us to estimate the basic reproduction number, R_0 , at an early stage of an outbreak and present some examples of the methodology. Before going into the details, we introduce the following identity equation, which is helpful in establishing a link between the concept in the previous and the current section:

$$\int_0^t \psi(\tau) \int_0^\tau f(t - \tau') d\tau' d\tau = \int_0^t \Psi(\tau) f(t - \tau) d\tau \quad (14)$$

Although the previous formalism is very general, in almost all practical situations, it is very difficult to obtain $\tilde{Z}(t - \tau, t)$, unless one is able to perform a very precise contact-tracing investigation. As a result, we prefer to approximate the excess degree per individual by the “expected excess degree”, i.e., $\bar{Z}(t - \tau, t) \simeq \frac{z_2}{z_1}$. It should be noted that this approximation is only valid when the following three conditions are met: a) the effect of degree correlation

between neighbors is negligible; b) the number of infected individuals is large enough to provide adequate sample size; and c) the number of infected individuals is small enough comparing to the network size. Since our objective is to calculate R_0 at an early stage, these assumptions remain completely plausible. It should also be noted that the expected excess degree at $t = 0$ is $\bar{Z}(0) = z_1$ and rapidly tends to $\frac{z_2}{z_1}$ as time progresses. This leads to a more precise equation $\bar{Z}(t - \tau, t) \simeq \left((\tilde{\mathcal{R}}(t) - 1) \frac{z_2}{z_1} + z_1 \right) / \tilde{\mathcal{R}}(t)$ which might be more suitable when $\tilde{\mathcal{R}}(t)$ is rather small. As before, we also assume that the time to removal t_R has the probability density $\psi(t_R)$. Replacing $\bar{Z}(t - \tau, t)$ by $\frac{z_2}{z_1}$ in (8), we obtain a new expression for the number of new infections at time t :

$$\tilde{J}(t) = \int_0^t \tilde{J}(t - \tau') \Psi(\tau) \tilde{R}_0(t, \tau) \phi(\tau, t_R \geq \tau) d\tau, \quad (15)$$

where $\tilde{R}_0(t, \tau) \equiv \tilde{T}(t, \tau, t_R \geq \tau) \frac{z_2}{z_1}$ is the only independent random variable.

The expected number of infectious and removed individuals are given by

$$\tilde{\mathcal{I}}(t) = \int_0^t \psi(\tau) \int_0^\tau \tilde{J}(t - \tau') d\tau' d\tau \quad (16)$$

and

$$\tilde{\mathcal{R}}(t) = \int_0^t \psi(\tau) \int_\tau^t \tilde{J}(t - \tau') d\tau' d\tau \quad (17)$$

Figures 7 and 8 show the number of infectious (left panel) and removed (right panel) individuals for a disease that spreads on binomial and exponential networks, respectively. Each panel shows three curves. The curve comprising of red circles corresponds to the computer simulation of an epidemic on the network; during the simulation the new case counts are recorded to create a synthetic “time series” for $J(t)$. The blue curve corresponds to the closed form equations (16) or (17) (for $\tilde{\mathcal{I}}(t)$ or $\tilde{\mathcal{R}}(t)$), and the green one corresponds to the scenarios when the identity in equation (14) is used. These figures show a perfect agreement between these three curves.




Figure 7: Number of infectious and removed individuals as a function of time for the binomial network ($z_1 = 5$, $\lambda_i = 0.12771$ and $\lambda_r = 0.25$).




Figure 8: Number of infectious and removed individuals as a function of time for the exponential network ($\kappa = 4$, $\lambda_i = 0.12771$ and $\lambda_r = 0.25$).

It should be noted that in the quasi-stochastic analysis, since $\tilde{Z}(t - \tau, t)$ is approximated by $\frac{z_2}{z_1}$, equation (12) takes the following simple form

$$\tilde{Z}^r(t) = \frac{z_2}{z_1} \tilde{\mathcal{R}}(t) \quad (18)$$

The expected transmissibility of removed individuals $\tilde{T}^r(t)$ can be obtained as an extension of Eq. (6) by replacing the removal rate distribution $\psi(t_R)$ with the conditional distribution of removal given it occurred before time t , defined as $\tilde{\psi}^r(t, t_R)$. The quantity $\tilde{\psi}^r(t, t_R)dt_R$ is proportional to the number of individuals already removed by time t that were removed after exactly t_R units of time, i.e., $\tilde{\psi}^r(t, t_R)dt_R \propto \psi(t_R) \int_0^{t-t_R} J(t')dt' dt_R$. This probability function, after incorporating the proper normalization, can be written as

$$\tilde{\psi}^r(t, t_R)dt_R = \psi(t_R) \frac{\int_0^{t-t_R} J(t')dt'}{\int_0^t \psi(t_R) \int_0^{t-t_R} J(t')dt' dt_R} dt_R \quad (19)$$

The expected transmissibility of removed individuals can then be calculated from

$$\tilde{T}^r(t) = \int_0^t \tilde{\psi}^r(t, t_R) T(t_R) dt_R \quad (20)$$

We now define the *reproduction number of the removed individuals* as

$$\tilde{R}_0^r(t) = \tilde{T}^r(t) \frac{z_2}{z_1}. \quad (21)$$

Combining the last equation with equations (13) and (18) one obtains that

$$\tilde{T}^r(t) \frac{z_2}{z_1} = \frac{\tilde{\mathcal{I}}^r(t) + \tilde{\mathcal{R}}^r(t)}{\tilde{\mathcal{R}}(t)}. \quad (22)$$

Now, considering the symmetry between different compartments in figure 6, it is straightforward to derive the following equation:

$$\begin{aligned} \tilde{R}_0^i &= \tilde{T}^i(t) \frac{z_2}{z_1} \\ &= \frac{\tilde{\mathcal{I}}^i(t) + \tilde{\mathcal{R}}^i(t)}{\tilde{\mathcal{I}}(t)} \end{aligned} \quad (23)$$

Using equation (22), the basic reproduction number $R_0 = T \frac{z_2}{z_1}$ can then be estimated as

$$\tilde{R}_0(t) = \frac{\tilde{\mathcal{I}}^r(t) + \tilde{\mathcal{R}}^r(t)}{\tilde{\mathcal{R}}(t)} \frac{T}{\tilde{T}^r(t)}. \quad (24)$$

On the right hand side of the last expression, the expected value of $\tilde{\mathcal{I}}^r(t) + \tilde{\mathcal{R}}^r(t)$ can be calculated as

$$\tilde{\mathcal{I}}^r(t) + \tilde{\mathcal{R}}^r(t) = \int_0^t \int_0^{t'} \tilde{J}(t') \eta(t, t', \tau) \zeta(t', \tau) d\tau dt' \quad (25)$$

where $\eta(t, t', \tau)$ is the fraction of infected individuals who are removed by time t and might have infected others at time $t' \leq t$; $\zeta(t', \tau)$ is the probability function that an infection at time t' was caused by any of these individuals. Let $\tilde{J}^i(t, \tau) \equiv \tilde{J}(t - \tau) \Psi(\tau)$, then $\eta(t, t', \tau)$ can be written as

$$\eta(t, t', \tau) = \frac{\tilde{J}^i(t', \tau) - \tilde{J}^i(t, \tau + t - t')}{\tilde{J}^i(t', \tau)} = \frac{\Psi(\tau) - \Psi(\tau + t - t')}{\Psi(\tau)} \quad (26)$$

Fig. 9 illustrates the concept. It shows how the new infection rate at time t' is due to the infectious rate $J^i(t', \tau')$ a fraction of whom $J^i(t, \tau)$ stays infectious at a later time t . It is straightforward to write $\zeta(t'; \tau)$ as

$$\zeta(t, \tau) = \frac{\tilde{J}^i(t, \tau) \frac{dT(\tau, t_R \geq \tau)}{d\tau}}{\int_0^t \tilde{J}^i(t, \tau') \frac{dT(\tau', t_R \geq \tau')}{d\tau'} d\tau'} \quad (27)$$




Figure 9: Dependency of the number of new infections at time t' on the number of infectious cases at time $t' - \tau$, $\tilde{J}^i(t', \tau)$.

Substituting the expressions of η and ζ in Eq. (25) we obtain

$$\tilde{\mathcal{I}}^r(t) + \tilde{\mathcal{R}}^r(t) = \int_0^t \tilde{J}(t') dt' \left[1 - \frac{\int_0^t \tilde{J}^i(t, \tau + t - t') \frac{dT(\tau, t_R \geq \tau)}{d\tau} d\tau}{\int_0^t \tilde{J}^i(t', \tau) \frac{dT(\tau, t_R \geq \tau)}{d\tau} d\tau} \right], \quad (28)$$

which only depends on case notification data and on the disease transmissibility. For a disease with constant removal rate λ_r , we have $\tilde{J}^i(t, \tau + t - t') = \tilde{J}(t' - \tau) \psi^c(\tau) \psi^c(t - t')$ and as a result $\tilde{\mathcal{I}}^r(t) + \tilde{\mathcal{R}}^r(t) = \tilde{\mathcal{R}}(t)$. Interestingly, in this case we also obtain $\tilde{\mathcal{I}}^i(t) + \tilde{\mathcal{R}}^i(t) = \tilde{\mathcal{I}}(t)$ and $\tilde{\mathcal{R}}^i(t) = \tilde{\mathcal{I}}^i(t)$. This means that the first fraction on the right hand side in Eq. (24) equals unity, and therefore $T^r(t) \frac{z_2}{z_1} = 1$. Then the expression for $R_0(t)$ takes the simpler form

$$R_0(t) = \frac{T}{\tilde{T}^r(t)}. \quad (29)$$




Figure 10: The expected reproduction number of removed individuals for the binomial (left panel; $z_1 = 5$, $\lambda_i = 0.12771$ and $\lambda_r = 0.25$) and exponential (right panel; $\kappa = 4$) networks in terms of the number of removed individuals.

5.1 Numerical examples

To test the framework presented so far, we performed epidemic spread simulations on the two networks introduced earlier (binomial and exponential) and in each case collected the “time series” of case counts resulted from the simulations. In Fig. 10 we present the reproduction number of the removed individuals, $R_0^r(t)$, as a function of the number of removed individuals, $\tilde{\mathcal{R}}(t)$. The symbols show the results from direct counting during the simulation, whereas the lines show the results obtained from evaluating analytically each of the terms in Eq. (22) based on the estimated infection rate. The green and red colors correspond to the left and right hand side of Eq. (22), respectively. The discrepancy at the very early stage is mainly due to approximating $\tilde{Z}(t)$ with $\frac{z_2}{z_1}$. The small asymptotic deviation of our estimate for the exponential network comes from finite size effects (Noel *et al.*, 2009).

Fig. 11 shows the expected reproduction number for the binomial (left panel) and the exponential (right panel) networks. The “real” values are $R_0 = 1.6864$ and 2.3749 , respectively.

It is worth noting that $T^r(t)$ is a functional of $\lambda_i(t)$, $\lambda_r(t)$ and $J(t)$ (see equations (20) and (19)). Therefore, when λ_r is constant, the condition $T^r(t)\frac{z_2}{z_1} = 1$ allows one to evaluate, from




Figure 11: The expected reproduction number for the binomial (left panel; $z_1 = 5$, $\lambda_i = 0.12771$ and $\lambda_r = 0.25$) and exponential (right panel; $\kappa = 4$) networks in terms of the number of removed individuals.




Figure 12: Estimates for the value of $\frac{z_2}{z_1}$ for the binomial (left) and exponential (right) networks, when λ_i and λ_r are known.

a given time series, one of the three quantities λ_i , λ_r or $\frac{z_2}{z_1}$, if the other two quantities are assumed to be known (this statement holds even if λ_r is a function of time, in which case a more elaborate equation (22) should be used). For instance, we simulated again the epidemic on the binomial and exponential networks presented earlier, but this time with constant values for λ_r and λ_i . Using these values and the obtained $\tilde{J}(t)$, we calculated the value of $\frac{z_2}{z_1}$, which is presented in figure (12) for the binomial (left panel) and exponential (right panel) networks. The green line shows our estimates using the above condition, while the red line is calculated directly from the network structure.




Figure 13: Estimates for the value of λ_i for the binomial (left) and exponential (right) networks, when $\frac{z_2}{z_1}$ and λ_r are known.

As a second example, we used the same epidemic simulations, but this time we used as input the values for λ_r and $\frac{z_2}{z_1}$ that were derived directly from simulations and estimated the value of λ_i which is shown in figure 13 for the binomial (left panel) and exponential (right panel) networks. Here again, one would observe very good agreement between the value of λ_i used in the simulations (black line) and its estimated value from the condition $T^r(t)\frac{z_2}{z_1} = 1$ (red circles).

6 Exponential regime and generation interval distribution

In this section we show the relationship between our results and the methodology developed by Wallinga and Lipsitch (Wallinga & Lipsitch, 2007), which is a special case of the framework presented in this paper. In the exponential regime, one can ignore the stochastic fluctuation of the quantities and replace them with their expected values, since the stochastic effects become much less pronounced. In particular, we can re-write equation (15) by ignoring the stochastic effects as following

$$J(t) = \int_0^t J(t - \tau) \Psi(\tau) R_0(\tau, t_R \geq \tau) \phi(\tau, t_R \geq \tau) d\tau \quad (30)$$

We define the expression inside the bracket as $\chi(\tau')$, which is the expected infection rate per infected individual. Equation (30) then takes the following simple form

$$J(t) = \int_0^t J(t - \tau') \chi(\tau') d\tau', \quad (31)$$

which is the well known Lotka renewal equation (Lotka, 1939; Kot, 2002).

From the definition of $\chi(\tau)$ it is clear that $\int_0^\infty \chi(\tau) d\tau = R_0$. In the exponential regime, after replacing $J(t) \approx \exp(-\alpha t)$, we obtain

$$1 = \int_0^\infty \exp(-\alpha \tau') \chi(\tau') d\tau' \quad (32)$$

It is worth mentioning that the exact exponential regime can be reached in the limit $t \rightarrow \infty$ for an infinite-size network and that is why the upper limit in the above integral is set to ∞ . This means that there should be a slight deviation from the exponential behavior for a “finite-size” system at “finite time” once the outbreak has surpassed the stochastic regime. Following the line of argument in Wallinga and Lipsitch (Wallinga & Lipsitch, 2007) we find that

$$\frac{1}{R_0} = \int_0^\infty \exp(-\alpha \tau) \hat{\chi}(\tau) d\tau, \quad (33)$$

where $\hat{\chi}(\tau) = \chi(\tau)/R_0$ is the *generation interval distribution*. This quantity relates the basic reproduction number to the Laplace transform of the generation interval distribution in the asymptotic case (infinite-size, infinite-time). It is straightforward to obtain the following equation

$$\hat{\chi}(\tau) = \frac{\Psi(\tau) \frac{dT(\tau, t_R \geq \tau)}{d\tau}}{\int_0^\infty \Psi(\tau') \frac{dT(\tau', t_R \geq \tau')}{d\tau'} d\tau'} \quad (34)$$

This equation describes the relationship between the generation interval distribution and the derivative of transmissibility with respect to the duration of infection.

Equation (32) has a simpler form for constant λ_r and λ_i . This can be obtained by replacing $\psi(\tau) = \lambda_r \exp(-\lambda_r \tau)$ and $T(\tau', \tau) = 1 - \exp(-\lambda_i \tau')$ inside equation (32)

$$1 = \frac{z_2}{z_1} \lambda_i \int_0^\infty \exp(-\tau \lambda_r) \exp(-\alpha \tau) \exp(-\lambda_i \tau) d\tau \quad (35)$$

$$= \frac{z_2}{z_1} \frac{\lambda_i}{\lambda_i + \lambda_r + \alpha} \quad (36)$$

or

$$\alpha = \left(\frac{z_2}{z_1} - 1 \right) \lambda_i - \lambda_r \quad (37)$$

This leads to $\alpha = 0.26084$ and $\alpha = 0.51626$ for the binomial and exponential networks, respectively, which are in excellent agreement with results shown in figures 3 and 4.

7 Conclusions

Using concepts from network theory, we presented a method which provides a reliable estimate of the basic reproduction number, R_0 , at an early stage of an outbreak. We provided the details of calculations and compared our results at each step against simulations. Case notification data (time series) is the main input to this analytical framework. As an outbreak begins to unfold, the pattern of spread depends substantially on the structure of the underlying contact network. This dependency, in fact, manifests itself in the formation of the time series of newly infected cases. The proposed methodology highlights the interplay between the heterogeneity in contacts (network structure), the estimates of the basic reproductive number and infection

transmissibility. Depending on the circumstances, this methodology can be used to infer other useful quantities as well. For infectious pathogens that cause repeated outbreaks, there is enough empirical evidence to establish the distribution of the duration of infectiousness as well as the recovery rate of individuals. In this case, in addition to the basic reproductive number, the proposed methodology can shed light on the structure of the underlying contact network by estimating the ratio $\frac{z_2}{z_1}$. This is an important piece of information, because in many circumstances it is not possible to capture and build a detailed contact network among individuals based on some network generative rules. The importance of this quantity becomes more apparent when an emerging infectious disease strikes a population. In this circumstance, there is much less information on the characteristics of the disease such as the duration of infectiousness and recovery rate, which in turn determine the transmissibility of disease. Knowledge of the transmissibility of a disease during an early stage of an epidemic can play a crucial role, as effective and cost-effective public health intervention strategies hinge on the degree of contagiousness of a disease. On the other hand, before the spread of disease becomes rampant, the structure of the contact network within a population remains more or less stable. Therefore, the estimated value of $\frac{z_2}{z_1}$ obtained during epidemic lulls from the time series corresponding to common infections can be used to estimate the transmissibility of an emerging infectious disease at an early stage of an emerging infectious outbreak. We demonstrated this concept by two examples. Our estimate for the reproduction number very quickly tends to the real value, thus enabling epidemiologists and policymakers to identify the optimal control strategies in real time.

8 Acknowledgments

BP would like to acknowledge the support of the Canadian Institutes of Health Research (grants no. MOP-81273 and PPR-79231), the Michael Smith Foundation for Health Research (Senior Scholar Funds) and the British Columbia Ministry of Health (Pandemic Preparedness Modeling Project). BD, JM and RM were supported by these grants.

References

- Abbey, H. 1952 An examination of the reed-frost theory of epidemics. *Human Biol*, **24**, 201–233.
- Addy, C. L., Longini, I. M. & Haber, M. 1991 A generalized stochastic model for the analysis of infectious disease final size data. *Biometric*, **47**, 961–974.
- Anderson, H. & Britton, T. 2000 *Stochastic Epidemic Models and their Statistical Analysis*. Springer: Berlin.
- Anderson, R. M. & May, R. M. 1991 *Infectious diseases of humans*. Oxford: Oxford University Press.
- Arino, J., Brauer, F., van den Driessche, P., Watmough, J. & Wu, J. 2007 A final size relation for epidemic models. *Math Biosci Eng*, **4**, 159–175.
- Ball, F. 1986 A unified approach to the distribution of total size and total area under the trajectory of infectives in epidemic models. *Adv Appl Prob*, **18**, 289–310.
- Ball, F. & Donnelly, P. 1995 Strong approximations for epidemic models. *Stoch Process Appl*, **55**, 1–21.
- Becker, N. 1976 Estimation for an epidemic model. *Biometric*, **32**, 769–777.

- Brauer, F. 2008 Age-of-infection and the final size relation. *Math Biosci Eng*, **5**, 681–690.
- Cauchemez, S., Boelle, P. Y., Donnelly, C. A., Ferguson, N. M., Thomas, G., Leung, G. M., Hedley, A. J., Anderson, R. M. & Valleron, A. J. 2006a Real-time estimates in early detection of SARS. *Emerging Infect Dis*, **12**, 110–113.
- Cauchemez, S., Boelle, P. Y., Thomas, G. & Valleron, A. J. 2006b Estimating in real time the efficacy of measures to control emerging communicable diseases. *Am J Epidemiol*, **164**, 591–597.
- Cox, D. R. & Oakes, D. 1984 *Analysis of survival data*. Chapman & Hall.
- Dietz, K. 1993 The estimation of the basic reproduction number for infectious diseases. *Stat Math Res*, **2**, 23–41.
- Fraser, C., Donnelly, C. A., Cauchemez, S., Hanage, W. P., Van Kerkhove, M. D., Hollingsworth, T. D., Griffin, J., Baggaley, R. F., Jenkins, H. E. *et al.* 2009 Pandemic potential of a strain of influenza A (H1N1): early findings. *Science*, **324**, 1557–1561.
- Gillespie, D. T. 2001 Approximate accelerated stochastic simulation of chemically reacting systems. *The Journal of Chemical Physics*, **115**(4), 1716–1733. (doi:10.1063/1.1378322)
- Gillespie, D. T. & Petzold, L. R. 2003 Improved leap-size selection for accelerated stochastic simulation. *The Journal of Chemical Physics*, **119**(16), 8229–8234. (doi:10.1063/1.1613254)
- Guttorp, P. 1991 *Statistical Inference for Branching Processes*. Wiley: New York.
- Hethcote, H. W. 2000 The Mathematics of Infectious Diseases. *SIAM Review*, **42**, 599–653.
- Higham, D. J. 2008 Modeling and simulating chemical reactions. *SIAM Review*, **50**(2), 347–368.

- Kermack, W. O. & McKendrick, A. G. 1927 A Contribution to the Mathematical Theory of Epidemics. *Proc Roy Soc Lond A*, **115**, 700–721.
- Kot, M. 2002 *Elements of Mathematical Ecology*. Cambridge University Pres.
- Lipsitch, M., Cohen, T., Cooper, B., Robins, J. M., Ma, S., James, L., Gopalakrishna, G., Chew, S. K., Tan, C. C. *et al.* 2003 Transmission dynamics and control of severe acute respiratory syndrome. *Science*, **300**, 1966–19670.
- Lotka, A. J. 1939 A contribution to the theory of self-renewing aggregates, with special reference to industrial replacement. *Ann of Math Stat*, **10**, 1–25.
- Ma, J. & Earn, D. J. 2006 Generality of the final size formula for an epidemic of a newly invading infectious disease. *Bulletin of Math Biol*, **68**, 679–702.
- Newman, M. E. J. 2002 Spread of epidemic disease on networks. *Phys Rev E*, **66**, 016128–016139.
- Noel, P. A., Davoudi, B., Dube, L. J., Brunham, R. C. & Pourbohloul, B. 2009 Time evolution of epidemic disease on finite and infinite networks. *Phys Rev E*, **79**, 026101–026115.
- Wallinga, J. & Lipsitch, M. 2007 How generation intervals shape the relationship between growth rates and reproductive numbers. *Proc. Biol. Sci.*, **274**, 599–604.
- Wallinga, J. & Teunis, P. 2004 Different epidemic curves for severe acute respiratory syndrome reveal similar impacts of control measures. *Am J Epidemiol*, **160**, 509–516.

A Appendix

To perform Monte Carlo simulations of epidemic propagation on a network, one first requires an explicit knowledge of that network structure. We have used the following method (Newman, 2002; Noel *et al.*, 2009) to produce a network related to given degree distribution: (i) Generate a random degree sequence k_j of length N subjected to the degree distribution p_k . (ii) Make sure that $\sum_j k_j$ is an even number since a link is composed of two "stubs." (iii) For each j , produce a node with k_j stubs. (iv) Randomly choose a pair of unconnected stubs and connect them together. Repeat until all unconnected stubs are exhausted. (v) Test for the presence of self-loops and repeated links. Remove the faulty stubs by randomly choosing a pair of connected stubs and rewire them to the former stubs. Repeat until no self-loops and/or repeated links are found.

To simulate the spread of disease on a contact network in continuous time we follow a Tau-Leaping approach (Gillespie, 2001; Gillespie & Petzold, 2003; Higham, 2008), which we describe below. The processes of disease transmission along one edge and the removal of infectious individuals are controlled by $\lambda_i(\tau)$ and $\lambda_r(\tau)$, respectively. We divide time into intervals of length Δt and ensure that $\lambda_i(\tau)\Delta t$ and $\lambda_r(\tau)\Delta t$ are small enough, such that the expected epidemic curve does not vary much by reducing Δt even further. At every Δt step, each infectious individuals recovers with probability $\lambda_r(\tau_j)\Delta t$, where τ_j is the age of infection of individual j . If an infectious individual does not recover, then it infects independently each of its susceptible contacts with probability $\lambda_i(\tau_j)\Delta t$.

